

$$2. (a) \quad n_{\text{HCl}} = \frac{3.05 \times 10^4 \text{ g}}{36.46 \text{ g/mol}}$$

$$n_{\text{HCl}} = 836.5 \text{ mol} \quad (\text{extra digits carried})$$

$$[\text{H}^+] = \frac{836.5 \text{ mol}}{806 \text{ L}}$$

$$[\text{H}^+] = 1.038 \text{ mol/L}$$

$$\text{pH} = -\log 1.038$$



$$\text{pH} = 0.016$$

$$\text{pOH} = 14 - 0.016$$

$$\text{pOH} = 13.984$$

The pH and pOH of the hydrochloric acid are 0.016 and 13.984, respectively.

- (b) Assumptions are that the temperature remains constant, and that water does not contribute a significant amount of $\text{H}^+_{(\text{aq})}$ or $\text{OH}^-_{(\text{aq})}$.



$$K_a = \frac{[\text{H}^+_{(\text{aq})}][\text{F}^-_{(\text{aq})}]}{[\text{HF}_{(\text{aq})}]}$$

ICE Table for the Dissociation of Hydrofluoric Acid			
	$\text{HF}_{(\text{aq})} \rightleftharpoons$	$\text{H}^+_{(\text{aq})}$	$+ \text{F}^-_{(\text{aq})}$
Initial concentration (mol/L)	2.0	0.0	0.0
Change in concentration (mol/L)	-x	+x	+x
Equilibrium concentration (mol/L)	2.0 - x	x	x

$$K_a = \frac{[\text{H}^+_{(\text{aq})}][\text{F}^-_{(\text{aq})}]}{[\text{HF}_{(\text{aq})}]}$$

$$K_a = 6.6 \times 10^{-4}$$

Predict whether a simplifying assumption is justified:

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = \frac{2.0}{6.6 \times 10^{-4}}$$

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = 3000$$

Since $3000 > 100$, we may assume that $2.0 - x \doteq 2.0$.

The equilibrium expression becomes

$$\frac{x^2}{2.00} \doteq 6.6 \times 10^{-4}$$

$$x \doteq 0.036$$

The 5% rule justifies the assumption that $2.0 - x \doteq 2.0$.

$$[\text{H}^+_{(\text{aq})}] = [\text{F}^-_{(\text{aq})}] = 0.036 \text{ mol/L}$$

The concentrations of hydrogen and fluoride in a 2.0 mol/L hydrofluoric acid solution are 0.036 mol/L.

$$5. \quad [\text{H}^+] = \frac{K_w}{[\text{OH}^-]} = \frac{1.0 \times 10^{-14}}{2.5 \times 10^{-7}}$$

$$[\text{H}^+] = 4.0 \times 10^{-8} \text{ mol/L}$$

$$\text{pH} = -\log 4.0 \times 10^{-8}$$

$$\text{pH} = 7.40$$

The hydrogen ion concentration and pH of blood are $4.0 \times 10^{-8} \text{ mol/L}$ and 7.40, respectively.

$$6. \quad n_{\text{HC}_2\text{H}_3\text{O}_2(\text{aq})} = \frac{60\,000 \text{ g}}{60.06 \text{ g/mol}}$$

$$n_{\text{HC}_2\text{H}_3\text{O}_2(\text{aq})} = 999 \text{ mol}$$

$$C_{\text{HC}_2\text{H}_3\text{O}_2(\text{aq})} = \frac{999 \text{ mol}}{1250 \text{ L}}$$

$$C_{\text{HC}_2\text{H}_3\text{O}_2(\text{aq})} = 0.799 \text{ mol/L}$$

ICE Table for the Dissociation of Acetic Acid			
	$\text{HC}_2\text{H}_3\text{O}_2(\text{aq}) \rightleftharpoons$	$\text{H}^+(\text{aq}) +$	$\text{C}_2\text{H}_3\text{O}_2^-(\text{aq})$
Initial concentration (mol/L)	0.799	0.0	0.0
Change in concentration (mol/L)	-x	+x	+x
Equilibrium concentration (mol/L)	0.799 - x	x	x

$$\frac{[\text{H}^+(\text{aq})][\text{C}_2\text{H}_3\text{O}_2^-(\text{aq})]}{[\text{HC}_2\text{H}_3\text{O}_2(\text{aq})]} = 1.8 \times 10^{-5}$$

$$\frac{x^2}{0.799 - x} = 1.8 \times 10^{-5}$$

Predicting whether $0.799 - x \doteq 0.799 - x \dots$

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = \frac{0.799 \text{ mol/L}}{1.8 \times 10^{-5}}$$

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = 44400$$

Since $44400 > 100$, we assume that $0.799 - x \doteq 0.799$

$$\frac{x^2}{0.799} \doteq 1.8 \times 10^{-5}$$

$$x = 1.8 \times 10^{-5} \quad (\text{extra digits carried})$$

The 5% rule justifies the assumption.

$$[\text{H}^+(\text{aq})] = 3.795 \times 10^{-3} \text{ mol/L}$$

$$\text{pH} = -\log [\text{H}^+(\text{aq})]$$

$$= -\log [3.795 \times 10^{-3}]$$

$$\text{pH} = 2.421$$

$$\text{pOH} = 14 - 2.421$$

$$\text{pOH} = 11.579$$

The pH and pOH of the acetic acid solution are 2.421 and 11.579.

7. (a)

ICE Table for the Ionization of ASA			
	$\text{HC}_{10}\text{H}_7\text{CO}_{4(\text{aq})} \rightleftharpoons$	$\text{H}^+_{(\text{aq})} +$	$\text{C}_{10}\text{H}_7\text{CO}_{4(\text{aq})}^-$
Initial concentration (mol/L)	0.018	0.0	0.0
Change in concentration (mol/L)	-x	+x	+x
Equilibrium concentration (mol/L)	0.018 - x	x	x

$$\frac{[\text{H}^+_{(\text{aq})}][\text{C}_{10}\text{H}_7\text{CO}_{4(\text{aq})}^-]}{[\text{HC}_{10}\text{H}_7\text{CO}_{4(\text{aq})}]} = 3.27 \times 10^{-4}$$

$$\frac{x^2}{0.018 - x} = 3.27 \times 10^{-4}$$

Predicting whether $0.018 - x \doteq 0.018$...

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = \frac{0.018 \text{ mol/L}}{3.27 \times 10^{-4}}$$

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = 55$$

Since $55 < 100$, we cannot assume that $0.018 - x \doteq 0.018$.

$$\frac{x^2}{0.018} \doteq 3.27 \times 10^{-4}$$

$$\frac{x^2}{0.018 - x} = 3.27 \times 10^{-4}$$

$$(0.018 - x)(3.27 \times 10^{-4}) = x^2$$

$$5.89 \times 10^{-6} - 3.27 \times 10^{-4}x = x^2$$

$$x^2 + 3.27 \times 10^{-4}x - 5.89 \times 10^{-6} = 0$$

$$x = \frac{-3.27 \times 10^{-4} \pm \sqrt{(3.27 \times 10^{-4})^2 - 4(1)(-5.89 \times 10^{-6})}}{2(1)}$$

$$= \frac{-3.27 \times 10^{-4} \pm 4.86 \times 10^{-3}}{2}$$

$$x = 2.27 \times 10^{-3} \text{ or } -8.13 \times 10^{-4}$$

Since negative concentrations are meaningless,

$$x = 2.27 \times 10^{-3}$$

$$[\text{H}^+_{(\text{aq})}] = 2.27 \times 10^{-3} \text{ mol/L}$$

$$\text{pH} = -\log[\text{H}^+_{(\text{aq})}]$$


$$\text{pH} = -\log[2.27 \times 10^{-3}]$$

$$\text{pH} = 2.644$$


The pH of the ASA solution is 2.644.

- (b) An increase in temperature shifts the autoionization equilibrium of water to the right, producing more $\text{H}^+_{(\text{aq})}$. As a result, the pH of the solution would decrease.

8. (a)



$$\frac{[\text{H}^+_{(\text{aq})}][\text{CN}^-_{(\text{aq})}]}{[\text{HCN}_{(\text{aq})}]} = 6.2 \times 10^{-10}$$



$$\frac{[\text{H}^+_{(\text{aq})}][\text{CN}^-_{(\text{aq})}]}{[\text{HCN}_{(\text{aq})}]} = 6.2 \times 10^{-10}$$

$$\frac{x^2}{0.10 - x} = 6.2 \times 10^{-10}$$

Predicting whether $0.10 - x \doteq 0.10 \dots$

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} = \frac{0.10 \text{ mol/L}}{6.2 \times 10^{-10}}$$

$$\frac{[\text{HA}]_{\text{initial}}}{K_a} \doteq 10^9$$

Since $10^9 > 100$, we assume that $0.10 - x \doteq 0.10$.

$$\frac{x^2}{(0.10)} \doteq 6.2 \times 10^{-10}$$

$$x = 7.9 \times 10^{-6}$$

The 5% rule justifies the assumption.

$$[\text{H}^+_{(\text{aq})}] = 7.9 \times 10^{-6} \text{ mol/L}$$

$$\text{pH} = -\log[\text{H}^+_{(\text{aq})}]$$

$$= -\log[7.9 \times 10^{-6}]$$

$$\text{pH} = 5.10$$

The pH of the hydrocyanic acid is 5.10.

Calculating percent ionization ...


$$[\text{H}^+_{(\text{aq})}] = \frac{p}{100} \times [\text{HA}_{(\text{aq})}]$$

$$p = \frac{[\text{H}^+_{(\text{aq})}]}{[\text{HA}_{(\text{aq})}]} \times 100\%$$

$$= \frac{7.9 \times 10^{-6} \text{ mol/L}}{0.100 \text{ mol/L}} \times 100\%$$

$$= 7.9 \times 10^{-3} \%$$

The percent ionization of hydrocyanic acid is $7.9 \times 10^{-3} \%$.



$$\text{pH} = 8.65$$

$$\text{pOH} = 5.35$$

$$[\text{OH}^-_{(\text{aq})}] = 10^{-5.35}$$

$$= 4.47 \times 10^{-6} \text{ mol/L (extra digits carried)}$$

$$[\text{OH}^-_{(\text{aq})}] = [\text{H}_2\text{C}_6\text{H}_6\text{O}_2_{(\text{aq})}]$$

ICE Table for the Ionization of Ascorbate				
	$\text{HC}_6\text{H}_6\text{O}_2^-_{(\text{aq})} +$	$\text{H}_2\text{O}_{(\text{l})} \rightleftharpoons$	$\text{H}_2\text{C}_6\text{H}_6\text{O}_2_{(\text{aq})} +$	$\text{OH}^-_{(\text{aq})}$
Initial concentration (mol/L)	0.15	–	0.00	0.00
Change in concentration (mol/L)	-4.47×10^{-6}	–	4.47×10^{-6}	4.47×10^{-6}
Equilibrium concentration (mol/L)	$0.15 - (4.47 \times 10^{-6})$	–	4.47×10^{-6}	4.47×10^{-6}

$$\frac{[\text{H}_2\text{C}_6\text{H}_6\text{O}_2(\text{aq})][\text{OH}^-(\text{aq})]}{[\text{HC}_6\text{H}_5\text{O}_2^-(\text{aq})]} = K_b$$

$$\frac{(4.47 \times 10^{-6})^2}{0.15 - 4.47 \times 10^{-6}} = K_b$$

Assume that $0.15 - 4.47 \times 10^{-6} \doteq 0.15$.

$$\frac{(4.47 \times 10^{-6})^2}{0.15} = 1.3 \times 10^{-10}$$


K_b for the ascorbate ion is 1.3×10^{-10} .

10. (a)

acid	conjugate base	acid	conjugate base
$(\text{C}_6\text{H}_5)_3\text{CH}$	$(\text{C}_6\text{H}_5)\text{C}^-$	$\text{C}_4\text{H}_4\text{NH}$	$\text{C}_4\text{H}_4\text{N}^-$
$\text{HC}_2\text{H}_3\text{O}_2$	$\text{C}_2\text{H}_3\text{O}_2^-$	H_2S	HS^-
$(\text{C}_6\text{H}_5)_3\text{CH}$	$(\text{C}_6\text{H}_5)\text{C}^-$	OH^-	O^{2-}
$\text{C}_4\text{H}_4\text{NH}$	$\text{C}_4\text{H}_4\text{N}^-$	H_2S	HS^-

(b) (strongest acid) $\text{HC}_2\text{H}_3\text{O}_2$; H_2S ; $\text{C}_4\text{H}_4\text{NH}$; $(\text{C}_6\text{H}_5)_3\text{CH}$ (weakest acid)

11. The final solution is basic because of the hydrolysis of the methanoate ion:



12.

Ionic			Molecular		
neutral	acidic	basic	neutral	acidic	basic
CaCl_2	NH_4Cl	NaOH	$\text{C}_6\text{H}_{12}\text{O}_6$	HCl	NH_3
NaNO_3	$\text{Al}(\text{NO}_3)_3$	Na_2CO_3	O_2	H_2S	N_2H_4


13. (a) yellow

(b) red

(c) yellow-green

(d) colourless

(e) yellow



25.0 mL 17.9 mL

C 1.5 mol/L

$$n_{\text{NaOH}(\text{aq})} = 17.9 \text{ mL} \times \frac{1.50 \text{ mol}}{1 \text{ L}}$$


$$n_{\text{NaOH}(\text{aq})} = 26.9 \text{ mmol}$$

$$n_{\text{H}_3\text{PO}_4(\text{aq})} = 26.9 \text{ mmol} \times \frac{1}{2} = 13.4 \text{ mmol}$$

$$C_{\text{H}_3\text{PO}_4(\text{aq})} = \frac{13.4 \text{ mmol}}{25.0 \text{ mL}}$$

$$C_{\text{H}_3\text{PO}_4(\text{aq})} = 0.537 \text{ mol/L}$$

The concentration of the phosphoric acid solution is 0.537 mol/L.



$$n_{\text{NH}_3(\text{aq})} = C_{\text{NH}_3(\text{aq})} \times V_{\text{NH}_3(\text{aq})}$$


$$= 0.10 \text{ mol/L} \times 10.0 \text{ mL}$$

$$n_{\text{NH}_3(\text{aq})} = 1.0 \text{ mmol}$$

Final solution volume = 20.0 mL

Entities remaining in solution at the equivalence point: $\text{Cl}^-_{(\text{aq})}$, $\text{NH}_4^+_{(\text{aq})}$, $\text{H}_2\text{O}_{(\text{l})}$


Since $\text{Cl}^-_{(\text{aq})}$ does not hydrolyze, the pH of the solution is determined by $\text{NH}_4^+_{(\text{aq})}$.



Since 1.0 mmol of $\text{NH}_3_{(\text{aq})}$ was present initially, 1.0 mmol of $\text{NH}_4^+_{(\text{aq})}$ is present at the equivalence point.

$$[\text{NH}_4^+_{(\text{aq})}] = \frac{1.0 \text{ mmol}}{20.0 \text{ mL}}$$

$$[\text{NH}_4^+_{(\text{aq})}] = 0.050 \text{ mol/L}$$



$$\frac{[\text{H}^+_{(\text{aq})}][\text{NH}_3_{(\text{aq})}]}{[\text{NH}_4^+_{(\text{aq})}]} = 5.8 \times 10^{-10}$$

ICE Table for the Hydrolysis of $\text{NH}_4^+_{(\text{aq})}$			
	$\text{NH}_4^+_{(\text{aq})}$	\rightleftharpoons	$\text{H}^+_{(\text{aq})} + \text{NH}_3_{(\text{aq})}$
Initial concentration (mol/L)	0.050		0.000
Change in concentration (mol/L)	-x		+x
Equilibrium concentration (mol/L)	0.050 - x		x



$$\frac{[\text{H}^+_{(\text{aq})}][\text{NH}_3_{(\text{aq})}]}{[\text{NH}_4^+_{(\text{aq})}]} = 5.8 \times 10^{-10}$$

$$\frac{x^2}{0.050 - x} = 5.8 \times 10^{-10}$$

Applying the hundred rule ...

$$\frac{[\text{NH}_4^+_{(\text{aq})}]_{\text{initial}}}{K_a} = \frac{0.050}{5.8 \times 10^{-10}} > 100$$

The equilibrium simplifies to

$$\frac{x^2}{0.050} = 5.8 \times 10^{-10}$$
$$x = 5.385 \times 10^{-6} \quad (\text{extra digits carried})$$

The 5% rule verifies the assumption.

$$[\text{H}^+_{(\text{aq})}] \doteq 5.385 \times 10^{-6} \text{ mol/L}$$

$$\text{pH} \doteq -\log [5.385 \times 10^{-6}]$$

$$\text{pH} = 5.27$$

The pH of the solution at the equivalence point is 5.27.

$$(b) \quad K_b = \frac{[\text{NH}_4^+_{(\text{aq})}][\text{OH}^-_{(\text{aq})}]}{[\text{NH}_3_{(\text{aq})}]}$$

$$K_b = 1.8 \times 10^{-5}$$

$$\frac{x^2}{0.10 - x} = 1.8 \times 10^{-5}$$

Predicting the validity of the assumption ...

$$\frac{0.10}{1.8 \times 10^{-5}} > 100$$

∴ we may assume that $0.10 - x \doteq 0.10$

The equilibrium expression becomes

$$\frac{x^2}{0.10} \doteq 1.8 \times 10^{-5}$$

$$x \doteq 1.342 \times 10^{-3} \quad (\text{extra digits carried})$$

The 5% rule verifies the simplification assumption.

$$[\text{OH}^-]_{(\text{aq})} = 1.342 \times 10^{-3} \text{ mol/L}$$

$$\text{pOH} = -\log[1.342 \times 10^{-3}]$$

$$\text{pOH} = 2.872$$

$$\text{pH} = 14 - \text{pOH}$$

$$= 14.0 - 2.872$$

$$\text{pH} = 11.13$$

The initial pH of the ammonia solution is 11.13.

(c) $V_{\text{HCl}(\text{aq})} = 5.00 \text{ mL}$

$$C_{\text{HCl}(\text{aq})} = 0.100 \text{ mol/L}$$

$$n_{\text{HCl}(\text{aq})} = V_{\text{HCl}(\text{aq})} \times C_{\text{HCl}(\text{aq})}$$

$$= 5.00 \text{ mL} \times 0.100 \text{ mol/L}$$

$$n_{\text{HCl}(\text{aq})} = 0.50 \text{ mmol}$$

$$V_{\text{NH}_3(\text{aq})} = 10.00 \text{ mL}$$

$$C_{\text{NH}_3(\text{aq})} = 0.100 \text{ mol/L}$$

$$n_{\text{NH}_3(\text{aq})} = V_{\text{NH}_3(\text{aq})} \times C_{\text{NH}_3(\text{aq})}$$

$$n_{\text{NH}_3(\text{aq})} = 10.00 \text{ mL} \times 0.100 \text{ mol/L}$$

$$n_{\text{NH}_3(\text{aq})} = 1.00 \text{ mmol}$$

$$n_{\text{NH}_3(\text{aq})} \text{ remaining ...}$$

$$n_{\text{NH}_3(\text{aq})} = 1.00 \text{ mmol} - 0.50 \text{ mmol}$$

$$n_{\text{NH}_3(\text{aq})} = 0.50 \text{ mmol}$$

$$\text{Total volume} = 5.00 \text{ mL} + 10.00 \text{ mL} = 15.00 \text{ mL}$$

$$C_{\text{NH}_3(\text{aq})} = \frac{0.50 \text{ mmol}}{15.00 \text{ mL}}$$

$$C_{\text{NH}_3(\text{aq})} = 0.0333 \text{ mol/L} \quad (\text{extra digits carried})$$

$$C_{\text{NH}_4^+(\text{aq})} = \frac{0.50 \text{ mmol}}{15.00 \text{ mL}}$$

$$C_{\text{NH}_4^+(\text{aq})} = 0.0333 \text{ mol/L} \quad (\text{extra digits carried})$$

Let x represent the changes in concentration that occur as the system reestablishes equilibrium.

ICE Table for the Ionization of Ammonia				
	$\text{NH}_3(\text{aq})$	$+ \text{H}_2\text{O}(\text{l}) \rightleftharpoons$	$\text{OH}^-(\text{aq}) +$	$\text{NH}_4^+(\text{aq})$
Initial concentration (mol/L)	0.0333	–	0.00	0.0333
Change in concentration (mol/L)	–x	–	+x	+x
Equilibrium concentration (mol/L)	$0.0333 - x$	–	x	$0.0333 + x$

$$\frac{[\text{NH}_4^+(\text{aq})][\text{OH}^-(\text{aq})]}{[\text{NH}_3(\text{aq})]} = K_b$$

$$\frac{x(0.0333 + x)}{0.0333 - x} = 1.8 \times 10^{-5}$$

Applying the hundred rule ...

$$\frac{[\text{NH}_3(\text{aq})]_{\text{initial}}}{K_a} = \frac{0.0333}{1.8 \times 10^{-5}}$$

$$\frac{[\text{NH}_3(\text{aq})]_{\text{initial}}}{K_a} = 1850$$

Since $1850 > 100$, we can assume that $0.0333 + x \doteq 0.0333$ and that $0.0333 - x \doteq 0.0333$

The equilibrium simplifies to

$$\frac{x(0.0333)}{0.0333} = 1.8 \times 10^{-5}$$

$$x = 1.8 \times 10^{-5}$$

The 5% rule validates the assumption.

$$[\text{OH}^-(\text{aq})] \doteq 1.8 \times 10^{-5} \text{ mol/L}$$

$$\text{pOH} \doteq -\log [1.8 \times 10^{-5}]$$

$$= 4.7445 \quad (\text{extra digits carried})$$

$$\text{pH} = 9.26$$

The pH after adding 5.0 mL of HCl is 9.26.

(d) Entities at the equivalence point are: $\text{H}_2\text{O}(\text{l})$, $\text{NH}_4^+(\text{aq})$, $\text{Cl}^-(\text{aq})$

(e) The equivalence point is reached after 10.0 mL of $\text{HCl}(\text{aq})$ is added. The pH of this solution is 5.27. (See calculation in (a).)

$$\begin{aligned} \text{(f)} \quad n_{\text{HCl}(\text{aq})} &= V_{\text{HCl}(\text{aq})} \times C_{\text{HCl}(\text{aq})} \\ &= 15.0 \text{ mL} \times 0.10 \text{ mol/L} \end{aligned}$$

$$n_{\text{HCl}(\text{aq})} = 1.5 \text{ mmol} \quad (\text{extra digits carried})$$

$$n_{\text{NH}_3(\text{aq})} = V_{\text{NH}_3(\text{aq})} \times C_{\text{NH}_3(\text{aq})}$$

$$\begin{aligned} n_{\text{NH}_3(\text{aq})} &= 10.0 \text{ mL} \times 0.10 \text{ mol/L} \\ &= 1.0 \text{ mmol} \end{aligned}$$

$$n_{\text{HCl}(\text{aq})} \text{ remaining} = 1.5 \text{ mmol} - 1.0 \text{ mmol}$$

$$= 0.5 \text{ mmol}$$

$$C_{\text{HCl}(\text{aq})} = \frac{n_{\text{HCl}(\text{aq})}}{V_{\text{HCl}(\text{aq})}}$$

$$= \frac{0.5 \text{ mmol}}{25 \text{ mL}}$$

$$C_{\text{HCl(aq)}} = 0.020 \text{ mol/L}$$

$$[\text{H}^+_{\text{(aq)}}] = 0.020 \text{ mol/L}$$


$$\text{pH} = -\log 0.020$$

$$\text{pH} = 1.70$$



After 15.0 mL of acid is added, the pH of the solution is 1.70.

- (g) A suitable indicator for this titration would be methyl red (pH range 4.4–6.2).
16. (a) NH_4^+
 (b) NH_2^-
 (c) $2 \text{NH}_3 \rightleftharpoons \text{NH}_4^+ + \text{NH}_2^-$
 (d)

Titration Curve for Titrating NH_2^- with NH_4^+



17. The pH would be between 5.0 and 5.4. An estimate of the hydrogen ion concentration is $1 \times 10^{-5} \text{ mol/L}$.



Applying Inquiry Skills

19. (i) Test the solutions with a pH meter. The strongest bases have the largest pH values.
 (ii) Test the solutions with universal indicator. Compare the indicator colour with a colour chart to determine the approximate pH.
20. (a) (i) neutral
 (ii) basic
 (iii) basic
 (iv) <7
 (v) <7
 (vi) basic
 (vii) <7
- (b) The predictions could be tested with a pH meter. (The meter should be carefully calibrated, and rinsed between solutions.)
21. (Sample answer—there are many possible correct solutions to this problem.) If the solutions are tested with a pH meter, and the pH values are ordered from smallest to largest, then the solutions are sulfuric acid, hydrochloric acid, acetic acid, ethanediol, ammonia, sodium hydroxide, and barium hydroxide, respectively.


Diagnostic Tests on the Unlabelled Solutions			
Litmus	Conductivity	Acid/Base titration	Analysis
red	low	one volume	CH ₃ COOH _(aq)
blue	very high	two volumes	Ba(OH) _{2(aq)}
blue	low	one volume	NH _{3(aq)}
no change	none	not applicable	C ₂ H ₄ (OH) _{2(aq)}
red	higher	two volumes	H ₂ SO _{4(aq)}
red	high	one volume	HCl _(aq)
blue	high	one volume	NaOH _(aq)

22. (a) There is no third pH endpoint and therefore, no third equivalence point in this titration.
 (b) Removing the hydroxide ions by precipitation will cause the equilibrium to shift to the right, producing more hydroxide ions.
 (c) Hydrochloric acid is not a primary standard.
 (d) Both reactants and products form basic solutions and litmus cannot be used to distinguish among basic solutions.
 (e) Cobalt chloride paper is used in a diagnostic test for the production of water in a strong acid–strong base reaction.
 (f) The strength of an acid is determined by titration.

Making Connections




- (b) Alkalosis is based on the carbonic acid/bicarbonate equilibrium:




During alkalosis, an increase in blood levels of the bicarbonate ion removes hydrogen ions, thereby increasing blood pH. Symptoms of alkalosis include vomiting, headache, and nausea.

- (c) Hydroxide-based antacids are very common.

Extension




Place an accurately measured volume of Ba(OH)_{2(aq)} into a beaker and immerse the leads from a conductivity meter in the solution. Measure and record the initial conductivity reading. Titrate with standardized H₂SO_{4(aq)} and record the conductivity readings. A sudden drop in the conductivity serves as the endpoint of the titration. This drop in conductivity corresponds to the removal of most of the ions through the production of water and insoluble barium sulfate. When the H₂SO_{4(aq)} is added in excess, the conductivity increases due to the ions present in the acid.



From the first ionization:

Since sulfuric acid is a strong acid $[\text{H}^+_{(aq)}] = 1.0 \times 10^{-2} \text{ mol/L}$

From the second ionization:



$$K_a = \frac{[\text{H}^+_{(aq)}][\text{HSO}_4^-_{(aq)}]}{[\text{HSO}_4^-_{(aq)}]}$$

$$K_a = 1.0 \times 10^{-2}$$

ICE Table for the Ionization of the Hydrogen Sulfate Ion			
	HSO ₄ ⁻ _(aq)	⇌	H ⁺ _(aq) + SO ₄ ²⁻ _(aq)
Initial concentration (mol/L)	0.010		0.00
Change in concentration (mol/L)	-x		+x
Equilibrium concentration (mol/L)	0.010 - x		x